

Arkansas Urology Update Form

Patient Name: _____ Date of Birth: _____

Address: _____

Doctor Sign Off

Phone: _____ Work Phone: _____

Cell Phone: _____

Primary Insurance Company: _____

ID#: _____ Group#: _____

Secondary Insurance Company: _____

ID#: _____ Group#: _____

Pharmacy name & phone: _____

Referring DR: _____ PCP: _____

Medications:

_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____

Medical Conditions:

_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____

Surgical History:

_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____

Allergies:

_____	_____	_____
_____	_____	_____
_____	_____	_____


Patient Signature: _____

IF YOU'RE A MAN

Take these tests and find out about 2 common conditions

Erection difficulties are when a man can't get or keep an erection. This is also known as erectile dysfunction (ED). About 30 million men in the US have some degree of ED. Also common in men is benign prostatic hyperplasia (BPH). This is a urinary problem in men older than 50.

It may feel awkward to talk to your doctor. But you are not alone. Many men have talked to their doctors about these problems. So take these tests. And help your doctor find out if treatment is right for you.



Arkansas Urology

Could you benefit from treatment for BPH?

Please circle the answer that *best describes* your response to each of the following questions.

Not at all	Less than 1 time in 5	Less than half the time	About half the time	More than half the time	Almost always
0	1	2	3	4	5

1. Incomplete emptying Over the past month, how often have you had a sensation of not emptying your bladder completely after you have finished urinating?

0	1	2	3	4	5
---	---	---	---	---	---

2. Frequency Over the past month, how often have you had to urinate again less than 2 hours after you have finished urinating?

0	1	2	3	4	5
---	---	---	---	---	---

3. Intermittency Over the past month, how often have you found you stopped and started again several times when you urinated?

0	1	2	3	4	5
---	---	---	---	---	---

4. Urgency Over the past month, how often have you found it difficult to postpone urination?

0	1	2	3	4	5
---	---	---	---	---	---

5. Weak stream Over the past month, how often have you had a weak urinary stream?

0	1	2	3	4	5
---	---	---	---	---	---

6. Straining Over the past month, how often have you had push or strain to begin urination?

0	1	2	3	4	5
---	---	---	---	---	---

7. Nocturia Over the past month, how many times did you most typically get up to urinate from the time you went to bed at night until the time you got up in the morning?

0	1	2	3	4	5+
---	---	---	---	---	----

Score _____

Add the numbers from your answers to questions 1 through 7. The maximum possible score is 35. Remember: This information is not intended as a substitute for medical treatment.

Note: This test is used to measure the severity of your symptoms. It is not a diagnostic test. In other words, it will not tell you whether or not you have BPH. Talk to your doctor to determine if your symptoms are due to BPH.

Adapted from Barry MJ, et al. The American Urological Association symptom index for benign prostatic hyperplasia. *J Urol.* 1992; 148:1549-1557.

Could you benefit from treatment for ED?

Each question has several possible responses. Please circle the number of the response that *best describes* your own situation. Make sure that you select only one response for each question.

Over the past 6 months:

1. How do you rate your confidence that you could get and keep an erection?

Very low	Low	Moderate	High	Very high
1	2	3	4	5

2. When you had erections with sexual stimulation, how often were your erections hard enough for penetration (entering your partner)?

No Sexual activity	Almost never or never	A few times (much less than half the time)	Sometimes (about half the time)	Most times (much more than half the time)	Almost always or always
0	1	2	3	4	5

3. During sexual intercourse, how often were you able to maintain your erection after you had penetrated (entered) your partner?

Did not attempt intercourse	Almost never or never	A few times (much less than half the time)	Sometimes (about half the time)	Most times (much more than half the time)	Almost always or always
0	1	2	3	4	5

4. During sexual intercourse, how difficult was it to maintain your erection to completion of intercourse?

Did not attempt intercourse	Extremely difficult	Very difficult	Difficult	Slightly difficult	Not difficult
0	1	2	3	4	5

5. When you attempted sexual intercourse, how often was it satisfactory to you?

Did not attempt intercourse	Almost never or never	A few times (much less than half the time)	Sometimes (about half the time)	Most times (much more than half the time)	Almost always or always
0	1	2	3	4	5

Score _____

Add the numbers corresponding to questions 1 through 5. If your score is 21 or less, you may want to talk to your doctor.

Adapted from Rosen RC, et al. Development and evaluation of an abridged, 5-item version of the International Index of Erectile Function (IIEF-5) as a diagnostic tool for erectile dysfunction. *Int J Impot Res.* 1999; 11:319-326.